SUNSHINE AGRICULTURE
e-Newsletter

ISSN (E): 2583 – 0821

Sun. Agri.:e- Newsletter, (2026) *6*(10), 111-116

Article ID: 698

Biofortification and Nutritional Quality in Vegetables: From Fundamental Principles to Sustainable Applications

V Madan Mohan Rao[1]
Y. Bramaramba[2]

[1]HOD, Department of Botany and Horticulture, Lecturer in Botany, Govt Degree College Chintalapudi, Eluru, Dist., AP
[2]S.A (BIOLOGY), ZPHS Pragadavaram, Eluru Dist., A.P

Open Access

*Corresponding Author
V Madan Mohan Rao*

Available online at
www.sunshineagriculture.vitalbiotech.org

Article History
Received: 13. 10.2026
Revised: 18. 10.2026
Accepted: 23. 10.2026

This article is published under the terms of the Creative Commons Attribution License 4.0.

INTRODUCTION

Vegetables are essential components of a balanced diet because they supply vitamins, minerals, dietary fibre, antioxidants and various bioactive compounds. Regular consumption of diverse vegetables contributes to normal growth, immune function, metabolic regulation and the maintenance of overall health. Carrots, spinach, amaranth, tomato, sweet pepper, broccoli, cabbage, peas and other vegetable crops contain different combinations of nutrients that contribute to dietary diversity. However, the nutritional quality of vegetables can vary substantially among species, cultivars, production environments and cultivation practices. Increasing vegetable production alone does not necessarily guarantee adequate micronutrient intake if the crops produced have low concentrations of essential nutrients or are not consumed in sufficient quantities.

Micronutrient deficiencies remain an important global nutrition challenge. Inadequate intake of iron, zinc, vitamin A and other essential nutrients can adversely affect human health, particularly among vulnerable populations. The causes of micronutrient deficiencies are complex and include limited dietary diversity, poverty, inadequate access to nutrient-rich foods, poor nutrient absorption and certain health conditions. Improving the nutritional quality of commonly consumed crops is therefore an important component of broader food and nutrition security strategies.

Biofortification refers to the development or production of food crops with enhanced concentrations of essential nutrients or improved nutrient bioavailability in their edible portions. Unlike conventional food fortification, which generally adds nutrients during food processing, biofortification aims to improve nutritional characteristics during crop growth and development. This approach can complement dietary diversification, conventional fortification, supplementation and nutrition education.

2. Fundamental Principles of Biofortification

The scientific basis of biofortification involves the interaction of plant genetics, soil nutrient availability, root absorption, nutrient transport, metabolism and the accumulation of nutrients in edible plant parts. Nutrient concentrations in vegetables are influenced by both the amount of a nutrient available to the plant and the plant's capacity to absorb, transport and store it. Consequently, increasing soil nutrient levels does not always produce a proportional increase in the nutritional quality of harvested vegetables.

2.1 Nutrient uptake and accumulation in vegetables

Plants absorb mineral nutrients primarily through their roots from soil solution or nutrient solutions. Essential mineral elements, including nitrogen, phosphorus, potassium, calcium, magnesium, iron and zinc, perform different physiological functions in plants. Their availability depends on soil pH, organic matter, moisture, temperature, microbial activity and interactions among nutrients. For example, iron and zinc may become less available in certain alkaline soils, whereas excessive concentrations of some nutrients can interfere with the absorption of others.

2.2 Nutritional quality and bioavailability

Nutritional quality is determined not only by the concentration of nutrients in food but also by the extent to which those nutrients remain available for digestion, absorption and utilization by the human body. Bioavailability refers to the proportion of a nutrient that is absorbed and becomes available for physiological functions. It is influenced by the chemical form of the nutrient, the food matrix, processing methods and interactions with other dietary components.

Iron illustrates the importance of bioavailability. Vegetables generally provide non-haem iron, whose absorption can be influenced by dietary components such as vitamin C, phytates and certain polyphenols. Vitamin C can improve non-haem iron absorption, whereas some compounds may reduce it under particular dietary conditions. Zinc bioavailability can also be affected by the presence of phytate and the composition of the overall diet.

Vegetables containing provitamin A carotenoids can contribute to vitamin A nutrition, but their effectiveness depends on carotenoid concentration, food preparation, fat intake and individual absorption. Cooking may reduce some heat-sensitive vitamins while improving the accessibility of certain carotenoids through the breakdown of plant tissues. Thus, the success of biofortification should be evaluated by considering nutrient concentration, retention during preparation and potential bioavailability rather than concentration alone.

3. Major Approaches to Vegetable Biofortification

Biofortification can be achieved through agronomic management, conventional breeding, molecular breeding and biotechnology. These approaches can be used individually or in combination, depending on the crop, nutrient of interest, available genetic resources and production environment.

3.1 Agronomic biofortification

Agronomic biofortification involves the management of soil, fertilizers, irrigation and other production practices to increase the concentration of essential nutrients in edible plant parts. It is often the most immediately accessible approach because it can be implemented using existing vegetable varieties and locally available agricultural technologies.

The application of zinc fertilizers, for example, may improve zinc concentration in

vegetables when soil or plant zinc availability is limiting. Iron fertilizers may help correct iron deficiency in suitable production systems, while selenium application can increase selenium accumulation in some crops when appropriate and safe rates are used. However, responses vary among species and cultivars, and excessive application may cause toxicity, nutrient imbalances or unsafe concentrations in edible tissues.

3.2 Conventional breeding and genetic biofortification

Genetic biofortification involves the development of vegetable varieties that naturally accumulate higher concentrations of target nutrients in edible tissues. Conventional breeding uses genetic variation available within cultivated varieties and compatible related germplasm to combine desirable nutritional traits with agronomic performance.

Carrot breeding provides opportunities to improve provitamin A carotenoid concentrations, while tomato breeding can target carotenoids, vitamin C and other beneficial compounds. Leafy vegetables can be selected for improved mineral composition, although the final nutritional outcome also depends on soil conditions and crop management. The breeding process must consider the relationship between nutrient traits and other characteristics, including yield, taste, texture, appearance and storage quality.

4. Biofortification Opportunities in Major Vegetable Crops

Different vegetable groups offer distinct opportunities for improving nutritional quality. The choice of target nutrient should reflect existing nutrient gaps, dietary patterns, crop biology and the feasibility of producing nutritionally improved varieties or applying appropriate agronomic practices.

Leafy vegetables such as spinach, amaranth and other edible greens are important sources of folate, carotenoids and several minerals. They may be suitable candidates for improving selected micronutrient concentrations because the leaves are consumed directly. However, mineral concentrations can be affected by soil contamination, irrigation-water quality and growing conditions. Safe production practices and appropriate washing and handling are therefore essential.

Root vegetables such as carrots and radishes provide opportunities to improve the nutritional composition of edible storage organs. Carrot breeding has particularly strong potential for increasing provitamin A carotenoids. Root vegetables may also respond to improved mineral management, although the effectiveness of biofortification depends on the ability of the plant to transport and accumulate the target nutrient in the edible root.

Fruit vegetables, including tomato, capsicum, brinjal and cucurbits, can contribute vitamins, minerals and antioxidant compounds. Tomatoes are important sources of carotenoids, including lycopene, while coloured capsicum varieties can supply vitamin C and various carotenoids. Breeding and crop management can improve selected nutritional characteristics, but the response should be evaluated alongside fruit yield, taste, colour and shelf life.

Table 1: Major approaches and applications of biofortification in vegetable crops

Vegetable crop or group	Important nutritional targets	Potential biofortification approach	Sustainable application and considerations
Spinach and amaranth	Iron, folate, carotenoids and other minerals	Variety selection, balanced micronutrient management and suitable soil improvement	Monitor soil and irrigation-water quality; assess nutrient bioavailability
Carrot	Provitamin A carotenoids	Conventional breeding and selection of carotenoid-rich varieties	Maintain root quality, yield and colour stability across environments
Tomato	Lycopene, carotenoids and vitamin C	Genetic improvement and balanced crop nutrition	Evaluate maturity, fruit quality, storage and processing effects
Capsicum	Vitamin C and carotenoids	Selection of coloured cultivars and suitable production management	Maintain colour, marketability and nutrient stability
Cabbage and broccoli	Vitamin C, folate and selected minerals	Genetic selection and optimized nutrient management	Avoid excessive fertilizer application and minimize postharvest losses
Peas and beans	Protein, iron, zinc and other minerals	Conventional breeding and balanced soil fertility management	Assess yield, seed or pod quality and nutrient bioavailability
Sweet potato	Provitamin A carotenoids in suitable orange-fleshed types	Varietal development and dissemination of nutrient-rich planting material	Promote locally adapted varieties and evaluate consumer preferences
Onion and garlic	Selected minerals and bioactive sulphur compounds	Cultivar selection and balanced nutrient management	Maintain bulb quality, storage life and crop-specific nutrient balance
Mixed vegetable systems	Multiple vitamins, minerals and bioactive compounds	Crop diversification, integrated nutrient management and improved varieties	Improve dietary diversity while conserving soil and water resources

5. Nutritional Benefits and Public Health Significance

Biofortified vegetables can contribute to improved dietary quality by increasing the availability of selected nutrients in commonly consumed foods. Their potential public health benefits depend on the nutrient concentration achieved, the quantity consumed, the frequency of consumption, nutrient bioavailability and the presence of other dietary sources. They should therefore be viewed as one component of a broader food-based strategy for improving nutrition.

Iron-rich vegetables may contribute to dietary iron intake, but their effectiveness in improving iron status depends on absorption, overall dietary composition and the severity and causes of deficiency. Combining plant foods containing iron with vitamin C-rich foods can improve the absorption of non-haem iron. Similarly, vegetables containing provitamin A

carotenoids can contribute to vitamin A intake, especially when they are consumed regularly as part of a diverse diet.

Zinc-rich vegetables may improve dietary zinc intake, although the contribution depends on crop concentration and consumption patterns. The benefits may be particularly relevant in diets with limited animal-source foods, but the actual nutritional effect must be evaluated within the broader dietary context. No single biofortified crop can supply all essential nutrients or replace the importance of a varied diet.

6. Nutrient Retention During Harvesting, Storage and Processing

The nutritional benefits of biofortified vegetables can be reduced if nutrients are lost during harvesting, transportation, storage or cooking. Fresh vegetables continue to undergo metabolic processes after harvest, and their nutritional composition may change with temperature, humidity, light exposure and storage duration. Vitamin C and folate can be particularly sensitive to handling and processing conditions, while some carotenoids may be affected by oxidation and heat.

Appropriate harvesting maturity is important because nutrient concentrations change during crop development. Harvesting too early or too late may affect nutritional composition, texture, flavour and storage life. Gentle handling, clean packaging and suitable temperature management can reduce physical damage and postharvest deterioration.

7. Challenges in Biofortification and Sustainable Implementation

Despite its potential, vegetable biofortification faces several scientific, agronomic and socioeconomic challenges. One major challenge is the interaction between nutritional traits and crop productivity. Some varieties with enhanced nutrient concentrations may not perform consistently under all environmental conditions. Breeding programmes must therefore combine nutritional improvement with yield stability, disease resistance, climate adaptation and consumer-preferred quality.

Another challenge is the variability of agronomic biofortification. Soil pH, organic matter, irrigation, rainfall and nutrient interactions can influence nutrient availability and plant accumulation. Fertilizer application that increases nutrient concentrations in one environment may produce limited responses in another. Excessive nutrient application can also increase costs, create environmental pollution and, in some cases, cause unsafe accumulation in edible tissues.

8. Future Prospects and Emerging Technologies

The future of vegetable biofortification will increasingly depend on integrating conventional plant breeding, molecular genetics, digital agriculture, soil science and human nutrition. Advances in genomics and molecular markers can help identify genes and genetic regions associated with mineral accumulation, carotenoid biosynthesis and vitamin metabolism. These tools can accelerate the development of improved varieties, particularly for traits that are difficult to measure accurately across large breeding populations.

Genomic selection and high-throughput phenotyping may improve the efficiency of selecting varieties with favourable combinations of nutritional quality, yield and stress tolerance. Gene editing may offer additional opportunities to investigate or modify nutrient-related pathways, subject to crop-specific technical feasibility, biosafety evaluation and applicable regulations. These technologies should be supported by conventional field testing to confirm that improved traits remain stable under real production conditions.

ISSN (E): 2583 – 0821

CONCLUSION

Biofortification is a promising strategy for improving the nutritional quality of vegetables and strengthening food and nutrition security. Through agronomic management, conventional breeding, molecular technologies and sustainable crop production, it is possible to increase selected vitamins, minerals and beneficial compounds in edible plant parts. The success of biofortification depends on understanding nutrient uptake and accumulation, improving bioavailability, maintaining crop productivity and ensuring nutrient retention during storage and cooking.

Sustainable applications require balanced fertilizer use, healthy soils, climate-adapted varieties, safe irrigation, efficient resource management and careful monitoring of nutritional outcomes. Nutrient concentration alone is not sufficient; biofortified vegetables must also be affordable, palatable, safe and widely accessible. Moreover, biofortification should complement dietary diversity, food fortification and other nutrition interventions rather than replace them.