SUNSHINE AGRICULTURE
e-Newsletter

ISSN (E): 2583 – 0821

Sun. Agri.:e- Newsletter, (2026) *6*(10), 49-52

Article ID: 684

Nanotechnology in Entomology: Tiny Tools for Smarter Pest Control

Shudhanshu Baliyan[1]
Vishvendra[2], Nikhil Kumar[2]
Lakshya Yadav[3]

[1]Department of Entomology
Govind Ballabh Pant University
of Agriculture and Technology
Pantnagar, Uttarakhand, 263145
[2]Department of Entomology
Sardar Vallabhbhai Patel
University of Agriculture
and Technology, Meerut
Uttar Pradesh, 250110
[3]Department of Entomology
Indian Veterinary Research
Institute, Bareilly
Uttar Pradesh, 243122

Open Access

*Corresponding Author
Shudhanshu Baliyan*

Available online at
www.sunshineagriculture.vitalbiotech.org

Article History
Received: 04.10.2026
Revised: 09.10.2026
Accepted: 14.10.2026

This article is published under the terms of the Creative Commons Attribution License 4.0.

INTRODUCTION

Think of a pesticide as a passenger. In a conventional spray, it travels freely in a liquid. In a nano-formulation, it travels in a designed vehicle a tiny droplet, capsule or particle built to keep it stable, deliver it to a useful location and release it at an appropriate rate.

Two broad approaches sit under this umbrella. In one, the nanomaterial itself acts as the active ingredient; in the other, the nanomaterial is a carrier for a conventional insecticide, plant extract, microbial protein or genetic material. Researchers have explored polymeric and lipid particles, silica, metals and metal oxides, nanocapsules and nanoemulsions (Athanassiou *et al.,* 2018).

Nano-structures can change water dispersibility, stability against sunlight and heat, release rate and the way a product adheres to a leaf or interacts with an insect. These changes can be useful, but the benefit depends on the formulation, the pest and the environment.

How Can Nanotechnology Help Entomology?

Nanotechnology can provide three main services in pest control: protection, delivery and timing. Fragile ingredients can be shielded from environmental degradation, active ingredients can be dispersed, retained on surfaces or delivered across biological barriers and controlled-release systems can regulate how quickly an ingredient becomes available. These functions underpin many nano-formulated insecticides, botanical products and microbial biopesticides (Athanassiou *et al.,* 2018; Wang *et al.,* 2022).

Delivery is especially difficult for RNA interference (RNAi). Double-stranded RNA is fragile and must reach insect tissues to silence an essential gene. In soybean aphid experiments, a spray combining dsRNA with a polymer-based nanocarrier and detergent improved penetration through the aphid body wall and produced mortality of up to about 78%. Because the study used soybean seedlings, field performance still needs to be demonstrated (Yan *et al.,* 2020).

These applications fit best within IPM rather than replacing it. The promise is better-protected products, more selective delivery and earlier warning, combined with monitoring, thresholds and other compatible tactics.

Biopesticides in a Protective Shell

Botanical and microbial products are attractive components of sustainable pest management, but many are fragile. Essential oils are volatile, while microbial insecticidal proteins can be damaged by environmental conditions. Nano-formulations are therefore being investigated as protective systems that improve stability and release.

Neem provides a useful example. Neem oil containing azadirachtin can be formulated as a nanoemulsion and laboratory bioassays have reported greater mortality of adult rice weevils and red flour beetles than with non-formulated neem oil. Essential-oil nanoemulsions also demonstrate how strongly performance can depend on the crop or commodity being protected. In one stored-grain study, a Carlina acaulis essential-oil nanoemulsion produced 94.4% mortality of rice weevil adults on barley after seven days, compared with about 43% on oats and 18% on maize (Kavallieratos *et al.,* 2023). The same formulation therefore did not perform uniformly across commodities.

Microbial products can also gain protection from nano-formulation. In laboratory tests, Bacillus thuringiensis Cry1Ab packaged in nanomaterial-stabilised emulsions degraded more slowly under ultraviolet exposure and maintained greater residual activity against European corn borer larvae. The tests used purified protein and harsh UVC conditions, so performance under natural sunlight and field spraying requires confirmation (Jalali *et al.,* 2023).


From "More Chemical" to "Smarter Delivery"

The central idea is not to make a pesticide more poisonous, but to make a given amount do more useful work. Better protection can reduce premature degradation; improved dispersion or adhesion can help more of the active ingredient reach the intended target; and controlled release can prolong activity. These features could permit lower effective doses in some systems, but dose reductions must be demonstrated experimentally rather than assumed.

Large evidence reviews give a balanced picture. A critical assessment of nanopesticides and nanofertilizers reported potential efficacy advantages but also highlighted weaknesses in nano-specific quality assurance and experimental controls (Kah *et al.,* 2018). A later analysis of 314 studies found that nanopesticides were, on average, more effective against target organisms and less toxic to non-target organisms than conventional analogues, but the advantage was smaller in field trials and some adverse effects remained poorly understood (Wang *et al.,* 2022).

Eyes on the Field: Nano-Sensors and Smart Monitoring

Good pest management starts with knowing what is present and when. Nanotechnology adds a chemical sensing dimension to monitoring. Researchers have coated microscopic silicon cantilevers tiny vibrating beams with nanocomposites such as polyaniline with silver or graphene oxide. When target molecules bind to the coating, the vibration frequency changes. Such systems have detected pheromones and defensive chemicals from stink bugs and have also detected signals associated with southern green stink bug adults on soybean plants (Dias *et al.,* 2024).



These are early demonstrations rather than ready-made farm tools. A practical device must work reliably under variable temperature, dust, humidity and field backgrounds, while remaining affordable and easy to connect with traps and decision rules.

Benefits but Also Questions

At their best, nano-formulations could reduce application losses, improve stability, provide controlled release and increase delivery efficiency. If lower doses can achieve the same control, environmental exposure may also fall. However, small does not automatically mean safe. Nanoparticles can behave differently from dissolved chemicals, and non-target effects need to be assessed carefully.

A review of 172 publications concluded that more information is needed to characterise how bees are exposed to nanotechnology-based pesticides before pollinator risks can be properly understood. Broader evaluations have also identified gaps in nano-specific quality assurance (Hooven *et al.,* 2019; Kah *et al.,* 2018). Important questions remain about persistence, movement, transformation and effects in soil and water. Regulatory assessment requires additional information on nanomaterial characterisation, exposure and hazard (EFSA Scientific Committee, 2021).

Other barriers include manufacturing cost, scale-up, field delivery and the translation of laboratory performance into real agricultural conditions. Most applications remain under research and need direct comparison with conventional formulations under realistic field conditions (Hofmann *et al.,* 2020).

What Does the Future Hold?

Imagine a smart pest-management chain: sensors and cameras flag an early pest build-up; monitoring data are combined with weather and crop stage; a decision rule determines whether action is necessary; and a tailored product is applied precisely. Follow-up monitoring then checks efficacy and whether beneficial organisms were spared. Research already provides several pieces of this vision through nanosensors, nanocarriers for RNA and protected biopesticides (Dias *et al.,* 2024; Yan *et al.,* 2020; Jalali *et al.,* 2023).

The next priorities are convincing field trials, non-target and pollinator safety studies, rugged low-cost sensors, better material characterisation and clearer regulatory guidance (EFSA Scientific Committee, 2021; Hofmann *et al.,* 2020).

Nanotechnology in entomology is not about making pesticides stronger. It is about making pest management more precise: the right amount, in the right form, in the right

place, at the right time. Whether these tiny tools deliver big benefits will depend on honest field trials, transparent safety science, sensible regulation and evidence. In pest management, precision rather than power may be the real revolution.

REFERENCES

Athanassiou, C. G., Kavallieratos, N. G., Benelli, G., Losic, D., Usha Rani, P., & Desneux, N. (2018). Nanoparticles for pest control: current status and future perspectives. *Journal of Pest Science*, *91*(1), 1-15.

Dias, D. A., Fernandes, I. A., Machado, E. P., Pedott, L., Blassioli-Moraes, M. C., Borges, M., ... & Steffens, C. (2024). Nanosensors for detecting volatile compounds in pest management: a focus on agricultural sustainability. *ACS Agricultural Science & Technology*, *4*(11), 1250-1259.

EFSA Scientific Committee. (2021). Guidance on risk assessment of nanomaterials to be applied in the food and feed chain: human and animal health. *Efsa Journal*, *19*(8), e06768.

Hofmann, T., Lowry, G. V., Ghoshal, S., Tufenkji, N., Brambilla, D., Dutcher, J. R., ... & Wilkinson, K. J. (2020). Technology readiness and overcoming barriers to sustainably implement nanotechnology-enabled plant agriculture. *Nature Food*, *1*(7), 416-425.

Hooven, L. A., Chakrabarti, P., Harper, B. J., Sagili, R. R., & Harper, S. L. (2019). Potential risk to pollinators from nanotechnology-based pesticides. *Molecules*, *24*(24), 4458.

Jalali, E., Bel, Y., Maghsoudi, S., Noroozian, E., & Escriche, B. (2023). Enhancing insecticidal efficacy of Bacillus thuringiensis Cry1Ab through pH-sensitive encapsulation. *Applied Microbiology and Biotechnology*, *107*(20), 6407-6419.

Kah, M., Kookana, R. S., Gogos, A., & Bucheli, T. D. (2018). A critical evaluation of nanopesticides and nanofertilizers against their conventional analogues. *Nature nanotechnology*, *13*(8), 677-684.

Kavallieratos, N. G., Bonacucina, G., Nika, E. P., Skourti, A., Georgakopoulou, S. K. C., Filintas, C. S., ... & Benelli, G. (2023). The type of grain counts: effectiveness of three essential oil-based nanoemulsions against Sitophilus oryzae. *Plants*, *12*(4), 813.

Wang, D., Saleh, N. B., Byro, A., Zepp, R., Sahle-Demessie, E., Luxton, T. P., ... & Su, C. (2022). Nano-enabled pesticides for sustainable agriculture and global food security. *Nature nanotechnology*, *17*(4), 347-360.

Yan, S., Qian, J., Cai, C., Ma, Z., Li, J., Yin, M., ... & Shen, J. (2020). Spray method application of transdermal dsRNA delivery system for efficient gene silencing and pest control on soybean aphid Aphis glycines. *Journal of Pest Science*, *93*(1), 449-459.