SUNSHINE AGRICULTURE
e-Newsletter

ISSN (E): 2583 – 0821

Sun. Agri.:e- Newsletter, (2026) *6*(10), 53-57

Article ID: 685

Role of Neem-Based Products in Sustainable Management of Insect Pests in Mungbean

Shivam Singh[1]
Pankaj Kumar[1]
Nandini Singh[1]
Chandan Singh[1]
Manshi Rohewal[2], Nikita[2]

[1]Research Scholar, Sardar Vallabhbhai Patel University of Agriculture & Technology Meerut – 250110 (U.P.)
[2]CCS University, Meerut

Open Access

*Corresponding Author
Shivam Singh*

Available online at
www.sunshineagriculture.vitalbiotech.org

Article History
Received: 04.10.2026
Revised: 09.10.2026
Accepted: 14.10.2026

This article is published under the terms of the Creative Commons Attribution License 4.0.

INTRODUCTION

Mungbean (*Vigna radiata* L.), commonly known as green gram, is an important short-duration pulse crop cultivated widely in India. It is valued for its high-quality protein, essential nutrients and ability to improve soil fertility through biological nitrogen fixation. Despite its importance, mungbean productivity is often affected by several insect pests from seedling establishment to pod formation and maturity. Major insect pests associated with mungbean include whitefly (*Bemisia tabaci*), aphids, thrips, jassids, pod borers and other pod-feeding insects. Among these, sucking pests are particularly important because they directly reduce plant vigour and, in the case of whitefly, can also contribute to the spread of yellow mosaic disease. Farmers often depend heavily on synthetic insecticides for pest control. Although insecticides can provide rapid suppression, their repeated and indiscriminate use may contribute to insecticide resistance, resurgence of pests, residues and adverse effects on natural enemies and other beneficial organisms. Therefore, there is a growing need for safer, economical and environmentally compatible pest-management options. Neem-based products are among the most promising botanical tools for sustainable insect pest management. Neem contains several biologically active compounds, particularly azadirachtin, which can interfere with feeding, development, reproduction and behaviour of many insect pests. When properly incorporated into an Integrated Pest Management (IPM) programme, neem products can help reduce pest pressure while conserving beneficial organisms.

Major Insect Pests of Mungbean

Mungbean can be attacked by several insect pests during different stages of crop growth.

1. Whitefly

Whitefly is one of the most important sucking pests of mungbean. Adults and nymphs suck sap from leaves, resulting in weakening of plants and reduction in photosynthetic activity. Honeydew excreted by whiteflies may encourage the development of sooty mould. More importantly, whitefly acts as a vector of mungbean yellow mosaic virus. Severe infection can result in yellowing of leaves, poor plant growth and substantial yield reduction. Early monitoring of whitefly populations is therefore essential.

2. Aphids

Aphids occur in colonies on tender shoots, young leaves and other succulent plant parts. They suck plant sap and may cause curling, yellowing and distortion of leaves. Heavy infestation can reduce plant vigour and affect flowering and pod development.

3. Thrips

Thrips are tiny insects that feed on tender plant tissues. Feeding damage may appear as silvery patches, distortion and drying of young leaves. Severe infestation can interfere with normal plant growth and reproductive development.

4. Pod Borers and Other Pod-Feeding Insects

Pod-feeding insects can cause direct damage to developing seeds. Larvae may bore into pods and feed on developing grains, resulting in damaged seeds and reduced marketable yield. Because different pests occur at different crop stages, regular field scouting is essential for timely management.

Why Neem-Based Products?

Neem (*Azadirachta indica* A. Juss.) is traditionally known for its insecticidal and medicinal properties. Neem-based products have attracted considerable attention in modern agriculture because they contain compounds with insecticidal, antifeedant, repellent and growth-regulating properties.

The most widely studied compound is **azadirachtin**. Unlike many conventional insecticides that primarily act through rapid toxicity, neem products can affect insects in several ways. This multiple mode of action makes neem particularly useful as a component of an IPM programme.

Neem products can be prepared or formulated in different ways, including:

- Neem seed kernel extract (NSKE)
- Neem oil
- Azadirachtin-based formulations
- Neem leaf preparations
- Neem cake

Among these, azadirachtin-based commercial formulations are commonly used in crop protection programmes, while neem seed kernel extracts are frequently used as a botanical option where locally prepared materials are appropriate.

How Neem Products Affect Insect Pests

Neem does not necessarily kill insects immediately. Its effectiveness often results from several biological effects.

Antifeedant action

Neem compounds can reduce feeding by insects. Reduced feeding is particularly important against sucking and chewing pests because continued feeding is responsible for much of the crop damage.

Repellent effect

Neem may make treated plant surfaces less attractive to some insect pests. This can reduce pest settling and feeding.

Growth-regulating effect

Azadirachtin can interfere with insect growth and development. In susceptible insects, it may disturb moulting and prevent normal development from one stage to another.

Effect on reproduction

Neem compounds may reduce egg laying and affect reproductive processes in some insect species. This can help suppress the pest population over time.

Effect on insect behaviour

Neem can influence feeding, settling, movement and oviposition behaviour. These effects may contribute to gradual reduction in pest pressure.

Neem-Based Products Useful in Mungbean

1. Neem Seed Kernel Extract

Neem seed kernel extract is a traditional botanical preparation that contains naturally occurring neem compounds. Properly prepared extracts can be used as part of a botanical pest-management programme.

The effectiveness of a homemade preparation can vary considerably depending on seed quality, preparation method, concentration and storage. Therefore, commercial formulations with standardized active ingredients may provide more consistent results.

2. Neem Oil

Neem oil contains several bioactive compounds and can act through repellent, antifeedant and other biological effects. It may be particularly useful against early infestations of sucking pests when applied with good spray coverage.

3. Azadirachtin-Based Formulations

Commercial azadirachtin formulations contain a standardized concentration of the active ingredient. They are generally more convenient for farmers because the formulation and recommended dose are provided on the product label. The exact dose and spray concentration should always be based on the registered product label and current recommendations for mungbean.

4. Neem Cake

Neem cake is primarily used as an organic soil amendment and has additional pest-management value in certain cropping systems. Its use can contribute to integrated crop and soil management, although it should not be considered a substitute for foliar pest management when damaging insect populations are already present.

Neem and Integrated Pest Management

Neem products should not be considered a stand-alone solution for every insect pest. Their greatest value is obtained when they are integrated with other appropriate pest-management practices.

A practical IPM programme for mungbean can include the following components:

Regular field scouting

Farmers should inspect mungbean fields regularly, particularly during periods favourable for pest development. The undersides of leaves should be examined carefully for whiteflies, aphids and other sucking pests.

Use of healthy and suitable varieties

Growing well-adapted varieties with good agronomic performance can help maintain crop vigour and reduce losses associated with pest attack.

Removal of severely infected plants

Where yellow mosaic disease is present, severely affected plants should be removed at an early stage according to local crop-protection recommendations. This can help reduce sources of infection and should be combined with management of the whitefly vector.

Conservation of natural enemies

Predators such as ladybird beetles, lacewings and spiders, along with parasitoids and other beneficial organisms, play an important role in naturally suppressing pest populations.

Unnecessary broad-spectrum insecticide applications can harm these beneficial organisms. Neem-based products, when appropriately selected and used, can provide an additional option within a conservation-oriented IPM programme.

ISSN (E): 2583 – 0821

Need-based chemical control

Synthetic insecticides should be used only when pest populations reach economically important levels and when recommended interventions are necessary. Selection of registered insecticides should be based on current local recommendations and product labels.

Practical Pest Management Schedule

Crop stage	Important pests	Recommended approach
Seedling stage	Aphids, thrips, whitefly	Regular scouting and early intervention
Vegetative stage	Whitefly, aphids, thrips, jassids	Neem-based products/IPM measures when required
Flowering stage	Thrips, aphids and other pests	Frequent monitoring; protect pollinators and natural enemies
Pod formation	Pod borers and other pod-feeding insects	Monitor pods and use need-based IPM
Seed development	Pod-feeding pests	Timely management based on pest incidence

Note: Product concentration, dose, waiting period and number of applications should always follow the current registered label and recommendations applicable to the crop and region.

How Farmers Can Get Better Results from Neem Products

The effectiveness of botanical products depends strongly on correct application.

1. **Start early**
   Neem products generally perform better when pest populations are low or at an early stage of infestation. Waiting until a severe outbreak has developed may result in inadequate control.
2. **Ensure good spray coverage**
   Whiteflies, aphids and other sucking pests are often found on the undersides of leaves. Proper spray coverage is therefore important.
3. **Follow the recommended dose**
   Using too little product may result in poor control, while excessive use does not necessarily provide better protection and can increase costs.
4. **4. Apply during suitable weather**
   Spraying should be avoided during strong winds or immediately before rainfall. Application should follow the product label and local agricultural recommendations.
5. **Repeat only when necessary**
   Neem-based products are generally better suited to preventive and early-stage management than emergency knockdown control. Reapplication should be based on pest pressure and label recommendations.

Advantages of Neem-Based Pest Management

Neem-based products offer several potential advantages:

- Derived from a plant-based source
- Useful component of IPM
- Multiple biological effects on insect pests
- Can reduce pest feeding and reproduction
- May help conserve beneficial organisms when appropriately used
- Suitable for integration with cultural and biological control
- Can contribute to reduced dependence on conventional insecticides
- Supports more sustainable crop-protection practices

However, "natural" does not automatically mean "risk-free." Neem products should still be used carefully and according to the recommended label directions.

Limitations and Precautions

Neem-based products also have limitations. Their effectiveness may vary with pest species, pest population, formulation, environmental conditions and application quality. They generally do not provide the rapid knockdown effect associated with some conventional insecticides.

Farmers should therefore avoid expecting immediate mortality after every application.

The following precautions are important:

- Use only registered and recommended products.
- Follow the label dose and application instructions.
- Do not increase the concentration unnecessarily.
- Avoid spraying during peak activity of pollinators.
- Avoid indiscriminate mixing of neem products with other pesticides.
- Check compatibility before tank mixing.
- Store commercial formulations as recommended on the label.
- Use protective clothing and follow all safety instructions during spraying.

Neem-Based Products as a Component of Sustainable Agriculture

The future of pest management lies not in completely replacing one group of pesticides with another, but in using different compatible tools strategically. Neem-based products can occupy an important position within this approach.

For example, a farmer can combine crop monitoring, resistant or tolerant varieties where available, field sanitation, conservation of natural enemies, botanical products, biological control and need-based chemical intervention. Such an approach can reduce unnecessary pesticide applications and improve the sustainability of mungbean production.

CONCLUSION

Mungbean is an important pulse crop, but insect pests can significantly reduce its productivity and quality. Whiteflies, aphids, thrips and pod-feeding insects are among the major pests requiring timely attention. Neem-based products, particularly azadirachtin formulations, neem oil and neem seed-based preparations, provide useful botanical tools for managing insect pests. Their antifeedant, repellent, growth-regulating and reproductive effects can help suppress pest populations, particularly when interventions are made early. The greatest benefits can be achieved when neem products are combined with regular field scouting, cultural practices, biological control, conservation of natural enemies and need-based use of recommended insecticides.

For successful mungbean production, the message is simple: monitor the crop regularly, identify the pest correctly, act at the right time and use neem as an important component of an integrated and sustainable pest-management strategy.

Key Message for Farmers

"Do not wait for a pest outbreak. Scout your mungbean crop regularly, identify the pest correctly and use neem-based products at the right time as part of an integrated pest-management programme."