SUNSHINE AGRICULTURE
e-Newsletter

ISSN (E): 2583 – 0821

Sun. Agri.:e- Newsletter, (2026) *6*(10), 81-87

Article ID: 691

Turning Poor-Quality Roughages into Nutritious Feed: Physical, Biological and Chemical Methods for Dairy Animals

Sanjay Howal[1]
Satish Dakore[2]
Sidharth Sabale[3]
Tarun Kumar Varun[4]

[1]Sanskaram College of Veterinary & Animal Sciences Sanskaram University, Jhajjar Haryana, India
[2]ICAR–National Dairy Research Institute, Karnal, Haryana, India
[3]Department of Animal Husbandry, Government of Maharashtra, India
[4]ICAR–National Dairy Research Institute, Karnal, Haryana, India

Open Access

*Corresponding Author
Sanjay Howal*

Available online at
www.sunshineagriculture.vitalbiotech.org

Article History
Received: 11.10.2026
Revised: 15.10.2026
Accepted: 19.10.2026

This article is published under the terms of the Creative Commons Attribution License 4.0.

INTRODUCTION

Feeding is one of the most important factors influencing milk production, reproductive performance and the overall health of dairy animals. In many dairy farming systems, particularly in tropical and subtropical regions, the availability of high-quality green fodder is seasonal and often insufficient to meet the nutritional requirements of dairy cattle and buffaloes. Consequently, farmers depend heavily on crop residues such as wheat straw, paddy straw, maize stover, sorghum stover and other fibrous materials.

Although these roughages are readily available and relatively inexpensive, their feeding value is often limited by low protein content, poor digestibility, high fibre content and low palatability. When fed as the major component of the diet without appropriate supplementation, they may not provide sufficient nutrients to support optimum milk production.

The good news is that low-quality roughages can be converted into more useful livestock feed through suitable physical, chemical and biological treatments. These methods improve the accessibility of nutrients, increase digestibility, reduce feed wastage and, in some cases, enhance voluntary feed intake. The appropriate technique depends on the type of roughage, availability of resources, farm size, cost and the technical knowledge of the farmer.

1. Understanding low-quality roughages

Roughages are bulky feeds containing substantial amounts of fibre. Low-quality roughages generally include mature crop residues and dry grasses that contain limited protein and energy and are poorly digested by ruminants.
Common examples include:

- Wheat straw and paddy straw
- Maize stover and sorghum stover
- Mature, weathered grasses
- Dry legume residues of variable quality
- Other crop residues with high lignin or silica content

The major nutritional limitations of these feeds are low crude protein, high structural fibre, limited availability of fermentable energy and restricted access of rumen microorganisms to the plant cell wall. Lignin, in particular, forms a barrier around cellulose and hemicellulose, reducing their breakdown in the rumen. Paddy straw may also contain substantial silica, which further limits its feeding value.

The objective of roughage improvement is not simply to make straw softer or more attractive. It is to improve the availability of nutrients to rumen microorganisms and the animal while maintaining feed safety and economic feasibility.

Various methods to improve low-quality roughages:



2. Physical methods of roughage improvement

Physical treatments modify the size, structure, moisture content or accessibility of plant material. They are generally easier to adopt than chemical treatments, although their nutritional benefits vary.

2.1 Chaffing and chopping

Chaffing is one of the simplest and most widely used methods of improving crop residues. Long straw is cut into shorter pieces using a chaff cutter.

Benefits:

- Reduces feed wastage during feeding.
- Makes mixing roughages with green fodder and concentrates easier.
- Improves handling and uniform distribution of feed.
- May improve intake when long, coarse material is difficult for animals to consume.

However, chopping alone does not substantially increase the chemical digestibility of straw. Excessively fine grinding may reduce effective fibre, encourage rapid passage through the digestive tract and interfere with normal rumination. A practical chop length should therefore support easy consumption while preserving adequate physically effective fibre in the complete diet.

2.2 Grinding and milling

Grinding reduces roughages to smaller particles, increasing their surface area and potentially improving access by rumen microorganisms. It can also help prepare crop residues for pellet production or complete feed blocks.

However, very fine grinding may increase dust, reduce chewing activity and alter rumen fermentation. Grinding is most useful when it forms part of a balanced feed-processing system rather than being treated as a complete solution to poor-quality straw.

2.3 Soaking and washing

Soaking dry roughages in clean water can soften the material and reduce dust. Washing may remove some soluble substances, soil and surface contaminants.

Nevertheless, soaking and washing do not reliably improve the digestibility of the major fibrous components. They may also cause losses of soluble nutrients, increase the risk of spoilage and require additional labour and water. Wet-treated material should be fed promptly unless it is preserved under appropriate conditions.

2.4 Steam treatment and pressure processing

Steam treatment exposes plant material to heat and moisture, sometimes under pressure. Under controlled conditions, it can alter the structure of plant cell walls and increase fibre accessibility.

More intensive processes, such as steam explosion, can disrupt the plant structure and improve digestibility. However, these technologies require specialised equipment, energy and careful process control. Excessive heat treatment may cause undesirable chemical changes, including reduced availability of protein.

Thus, steam treatment is generally more relevant to organised feed-processing units than to small dairy farms.

3. Chemical methods of roughage improvement

Chemical treatments are particularly important because they can change the structure of plant cell walls rather than merely changing particle size. Alkali treatments can disrupt linkages between lignin and carbohydrates, swell the fibre and improve access by rumen microorganisms.

3.1 Urea treatment

Urea treatment is one of the most practical methods for improving cereal straws in many smallholder dairy systems. Under suitable conditions, urea is converted through urease activity into ammonia, which acts on the fibrous material. The treatment can improve digestibility and also increase the nitrogen content of the treated straw.

A commonly described approach for cereal straw uses approximately 4 kg of urea dissolved in enough water to provide around 40 litres of solution per 100 kg of straw. The solution is distributed uniformly through the straw, which is then compacted and covered in an airtight manner. The material is left to react for several weeks, often around three to four weeks under warm conditions, with longer periods potentially required in cooler weather.

The exact procedure must be adapted to the roughage, ambient temperature, moisture content and locally validated recommendations.

Advantages of urea treatment

- Can improve the digestibility and intake of cereal straw.
- Increases its nitrogen content.
- Can reduce the amount of untreated straw wasted.
- May improve the utilisation of crop residues when used in a balanced feeding programme.

Important precautions

Urea-treated straw must not be confused with simply adding urea directly to an animal's

feed. Incorrect mixing, excessive application or improper feeding can cause dangerous ammonia exposure and urea toxicity. Treatment should follow an established local protocol, and the straw should be adequately aerated after opening if residual ammonia is present. It must not be fed while strong residual ammonia is evident.

Urea treatment also does not provide sufficient true protein or energy to meet all the requirements of a high-producing dairy animal. A suitable source of fermentable energy, adequate protein, minerals and vitamins may still be necessary.

3.2 Anhydrous ammonia treatment

Anhydrous ammonia can be applied to dry crop residues under controlled conditions. It reacts with the plant material and may increase nitrogen content, alter fibre structure and improve digestibility.

Although effective, this technique requires specialised storage and application equipment. Anhydrous ammonia is hazardous, corrosive to certain materials and capable of causing severe chemical burns and respiratory injury. It should be handled only by trained personnel using appropriate engineering controls and protective equipment.

For these reasons, it is generally unsuitable for untrained farmers and should be considered only where safe infrastructure and expert supervision are available.

3.3 Sodium hydroxide treatment

Sodium hydroxide, or caustic soda, is a strong alkali that can break or loosen certain linkages in plant cell walls. Treatment may increase the digestibility of cellulose and hemicellulose and improve the utilisation of straw.

However, sodium hydroxide is highly corrosive. Incorrect treatment can leave excessive alkali in the feed, and poorly managed effluent can harm soil and water. Sodium levels in the diet must also be considered.

Although this method has demonstrated nutritional benefits under controlled conditions, it is generally less convenient for small-scale dairy farmers than safer, locally established alternatives.

3.4 Calcium hydroxide treatment

Calcium hydroxide, commonly known as hydrated lime, can be used as an alkaline treatment for selected crop residues. It can improve fibre accessibility, although the response depends on the crop material, treatment conditions, moisture and duration.

Its effectiveness may be lower or slower than that of some other alkalis. Excess calcium and poor mixing must also be avoided. Farmers should use only a validated feed-treatment protocol and should not assume that ordinary agricultural lime, quicklime and hydrated lime are interchangeable.

3.5 Alkali treatment: what should farmers remember?

Chemical treatment can improve the utilisation of roughage, but it is not a substitute for balanced nutrition. The treated material must be prepared uniformly, stored correctly and evaluated for feeding safety. Farmers should seek advice from a veterinary university, animal husbandry department or qualified livestock nutritionist before adopting a new chemical treatment.

Never mix chemicals experimentally into feed or feed animals material that has been treated using an unknown concentration or procedure.

4. Biological methods of roughage improvement

Biological methods use microorganisms, enzymes or fermentation to modify plant material. They can be attractive because they may reduce dependence on strong chemicals, but their effectiveness depends heavily on the biological agent and treatment conditions.

4.1 Fungal treatment

Certain fungi, particularly selected white-rot fungi, can degrade lignin and improve the accessibility of cellulose and hemicellulose. Research has investigated their use for improving straw and other lignocellulosic residues.

However, the process requires careful selection of fungal strains, suitable moisture and temperature, sufficient treatment time and control of contamination. Some fungi also consume valuable carbohydrates, reducing the energy content of the feed. Others may produce harmful metabolites or reduce palatability.

Consequently, fungal treatment should be used only with validated, feed-safe strains and established procedures. Uncontrolled fungal growth on stored straw is not a safe or acceptable substitute for deliberate biological treatment.

4.2 Enzyme treatment

Commercial cellulases, xylanases and related enzymes can help break down selected components of plant cell walls. They may be applied before feeding or incorporated into specific feed-processing systems.

The benefits depend on enzyme type, dose, application method, temperature, pH and the characteristics of the roughage. Some enzyme products improve fibre digestion under particular conditions, whereas others produce little measurable response.

Farmers should choose products specifically intended for livestock feed and follow the manufacturer's instructions. Enzymes should not be expected to compensate for severe deficiencies in dietary protein, energy or minerals.

4.3 Fermentation and silage making

Ensiling is a biological preservation method in which suitable microorganisms ferment plant sugars under anaerobic conditions, producing acids that help preserve the feed.

It is particularly useful for green fodder and sufficiently moist, fermentable crop materials. Very dry straw usually contains too little moisture and too little readily fermentable sugar for reliable silage making without suitable preparation or additives.

Where appropriate, chopping, moisture adjustment, compacting and airtight sealing can support good fermentation. Molasses or other suitable additives may sometimes be used to provide fermentable sugars, depending on the crop material and the recommended process.

Good silage should have a pleasant acidic smell and no visible mould. Slimy material, putrid odours or extensive mould growth indicate poor preservation, and the affected feed should not be offered to dairy animals.

4.4 Microbial inoculants

Selected lactic acid bacteria are commonly used as silage inoculants to encourage desirable fermentation and suppress undesirable microorganisms. Certain microbial products have also been investigated for their effects on fibre degradation.

Their benefits are not universal: they depend on crop composition, moisture, temperature and the microbial strains used. A silage inoculant should not automatically be regarded as a proven treatment for increasing the digestibility of dry straw.

4.5 Fermented total mixed ration

A total mixed ration (TMR) combines roughage, concentrates, protein sources, minerals and other ingredients in a uniform mixture. In some systems, controlled fermentation of the mixture can improve preservation and feeding convenience.

The main nutritional advantage of a well-formulated TMR is that it helps provide a more consistent and balanced supply of nutrients. Fermentation may offer additional benefits under appropriate conditions, but it cannot correct a poorly formulated ration by itself.

5. Comparison of different methods

Method	Main benefit	Major limitation	Suitability
Chopping	Easier handling and less wastage	Limited direct improvement in digestibility	Most dairy farms
Grinding	Smaller particles and easier mixing	Excessive fineness can impair rumination	Feed-processing systems
Steam treatment	Changes plant structure	Equipment and energy costs	Industrial or specialised units
Urea treatment	May improve digestibility and nitrogen content	Requires strict safety and process control	Farms with trained guidance
Ammonia treatment	Can improve fibre utilisation	Serious chemical hazards	Specialised facilities
Alkali treatment	Disrupts some cell-wall linkages	Corrosive chemicals and residue management	Controlled operations
Fungal treatment	Potential lignin degradation	Slow process and contamination risk	Research or validated systems
Enzyme treatment	Targets specific fibre components	Variable response and product cost	Where efficacy is established
Ensiling	Preserves suitable moist fodder	Requires adequate moisture and airtight storage	Farms with suitable green fodder
TMR preparation	Improves ration consistency	Requires accurate formulation and mixing	Small to large dairy farms

6. How to select the right method for dairy animals

The best method is not necessarily the most technologically advanced one. It is the one that improves nutrient utilisation safely and economically under local conditions.

For small dairy farms, a practical strategy may include the following:

1. **Use a chaff cutter.** Prepare straw at a suitable length and reduce wastage during feeding.
2. **Improve the overall ration.** Combine roughage with appropriate green fodder, energy sources, protein supplements and mineral mixture according to the animal's requirements.
3. **Consider urea treatment where appropriate.** Adopt it only when a reliable protocol, trained guidance and suitable storage conditions are available.
4. **Preserve good-quality fodder.** Use proper silage-making techniques when suitable green fodder is available in surplus.
5. **Avoid unverified biological or chemical products.** Seek evidence of safety and effectiveness before purchasing them.
6. **Monitor animal performance.** Observe feed intake, rumination, dung consistency, body condition and milk yield, and obtain veterinary advice if problems arise.

High-yielding cows and buffaloes require particularly careful ration formulation because their nutrient demands may exceed what treated straw can supply. Young calves also have different nutritional needs and should not be managed using the same feeding strategy as mature lactating animals.

7. Economic and environmental importance

Improving low-quality roughages can help reduce dependence on expensive purchased feeds, improve the utilisation of locally available crop residues and reduce wastage. It can also help farmers make better use of seasonal fodder resources.

Nevertheless, the economic benefit depends on treatment costs, labour, equipment, changes in feed intake, digestibility and milk production. A method that improves laboratory digestibility may not necessarily deliver a profitable increase in milk yield on every farm.

Farmers should therefore compare the cost per unit of usable feed and the animal's overall performance rather than judging a method solely by changes in the appearance or nitrogen content of the straw.

Better use of crop residues can also reduce the need to burn them in the field, provided collection, storage and transport are practical. This offers a potential environmental benefit alongside improved livestock feeding.

CONCLUSION

Low-quality roughages need not remain a low-value resource in dairy farming. Physical methods such as chopping improve handling and reduce wastage; chemical treatments such as urea treatment can improve fibre utilisation under controlled conditions; and biological approaches such as ensiling, selected enzyme applications and validated fungal treatments offer additional opportunities.

However, roughage treatment is only one part of successful dairy nutrition. No treatment can replace a balanced ration that supplies adequate energy, protein, minerals, vitamins and physically effective fibre. Safety, cost, local availability and correct application should guide every decision.

For many smallholder dairy farmers, the most practical approach is to combine efficient straw preparation, balanced supplementation, good fodder conservation and expert-guided treatment where justified. With these measures, crop residues can become a more useful component of sustainable and economical dairy production.